

Mental disorders and long-term labour market outcomes: nationwide cohort study of 2 055 720 individuals

Hakulinen C, Elovainio M, Arffman M, Lumme S, Pirkola S, Manderbacka K, Keskimäki I, Böckerman P. Mental disorders and long-term labour market outcomes: nationwide cohort study of 2 055 720 individuals.

Objective: To examine the associations between an onset of serious mental disorders before the age of 25 with subsequent employment, income and education outcomes.

Methods: Nationwide cohort study including individuals ($n = 2\,055\,720$) living in Finland between 1988–2015, who were alive at the end of the year they turned 25. Mental disorder diagnosis between ages 15 and 25 was used as the exposure. The level of education, employment status, annual wage or self-employment earnings, and annual total income between ages 25 and 52 (measurement years 1988–2015) were used as the outcomes.

Results: All serious mental disorders were associated with increased risk of not being employed and not having any secondary or higher education between ages 25 and 52. The earnings for individuals with serious mental disorders were considerably low, and the annual median total income remained rather stable between ages 25 and 52 for most of the mental disorder groups.

Conclusions: Serious mental disorders are associated with low employment rates and poor educational outcomes, leading to a substantial loss of total earnings over the life course.

C. Hakulinen^{1,2} ,
M. Elovainio^{1,2}, M. Arffman²,
S. Lumme², S. Pirkola^{3,4},
I. Keskimäki^{2,3}, K. Manderbacka²,
P. Böckerman^{5,6,7} 

¹Department of Psychology and Logopedics, Faculty of Medicine, University of Helsinki, Helsinki, ²National Institute for Health and Welfare, Helsinki, ³Faculty of Social Sciences, University of Tampere, Tampere, ⁴Pirkanmaa Hospital District, Tampere, ⁵School of Business and Economics, University of Jyväskylä, Jyväskylä, ⁶Labour Institute for Economic Research, Helsinki, Finland and ⁷IZA Institute of Labor Economics, Bonn, Germany

Key words: mental disorders; income; education; employment

Christian Hakulinen, University of Helsinki, PO Box 21, 00014 Helsinki, Finland. E-mail: christian.hakulinen@helsinki.fi

Accepted for publication June 24, 2019

Significant outcomes

- Serious mental disorders are associated with higher rates of not being employed and not having any secondary or higher education.
- Annual median earnings for the population without serious mental disorders increased from 15 000 € to 30 000 € between ages 25 and 52. On the contrary, annual median earnings were under 6000 € for most of the serious mental disorder groups.
- The median total income varied between 8000 € to 15 000 € for most serious mental disorder groups, indicating that income transfers compensate partly labour market losses due to serious mental disorders.

Limitations

- Only individuals who were treated in Finnish hospitals were examined in the current study, and the severity of mental disorders was not measured. Individuals who did not seek treatment or who received treatment in primary care or in outpatient clinics were included in the comparison population.
- The findings are not applicable to individuals with less severe mental disorders.

Introduction

Mental disorders are a leading cause of global disease burden. They lead to a loss of billions of dollars annually (1). From the specific mental disorders, substance abuse, psychotic, depressive and anxiety disorders are among the largest contributors to loss of disability-adjusted life years (2). The global cost of mental disorders is enormous—economic evaluations have indicated that the costs were approximately 2.5 trillion US dollars in 2010 and they are projected to increase by around 6 trillion US dollars in 2030 (3). Whereas direct costs—which refer to the costs associated with diagnosis and treatment—are considerable, indirect costs—referring to the ‘invisible’ costs such as unemployment and income losses—are typically the largest components of the total economic burden caused by mental disorders (3–7). For example, from the indirect costs associated with schizophrenia, unemployment amounted to around third of the total costs (7, 8).

Cross-sectional and longitudinal studies have shown that substance abuse disorders, schizophrenia and other psychoses, depression and anxiety disorders are each significantly associated with higher unemployment and lower income over time (9–20). It also seems that these associations are stronger for serious mental disorders (21, 22). However, considerable variation in employment rates between different mental disorders has been reported. Whereas 10–20% of individuals with schizophrenia have been reported to be employed (23, 24), the numbers have been around 60% among individuals with bipolar disorder (25) or a major depressive disorder (4). In addition, studies that have examined the association between mental disorders with educational attainment have produced partially mixed findings. Whereas cross-sectional and longitudinal survey studies have found that most mental disorders are associated with poor educational attainment (26, 27), a recent register-based study from the Netherlands found that schizophrenia, but not depression or bipolar disorder, were associated with lower likelihood of obtaining secondary or higher education (28). Taken together, it seems that mental disorders can lead to irregular work careers and permanently low income, thus increasing the total economic burden of mental disorders (29, 30).

However, large scale studies on the topic are still limited and, most studies have focused on the role of socioeconomic factors in developing mental disorders (31–33). Moreover, most studies have not examined whether serious mental disorders lead to a permanently poor socioeconomic position and to

what extent. This issue is of high policy importance as most mental disorders emerge before the age of 25 (34, 35), increasing substantially the risk of poor economic outcomes over life course. Furthermore, most previous studies have mainly compared schizophrenia with mood disorders (16, 28), leaving out substance abuse and anxiety disorders.

Aims of the study

We examined in a comprehensive national cohort study the association between serious mental disorders with subsequent employment, earnings, total income and educational outcomes from a work-life course perspective. To accomplish this, we combined national register-based information on mental disorders with comprehensive administrative register data on long-term labour market outcomes.

Methods

Study population

Our study population included the total working-age resident population living in Finland between 1988 and 2015, who were alive at the end of the year they turned 25. The individuals included in the current study were born between 1963 and 1990. The data were compiled through linking several registries: Finnish Longitudinal Employer–Employee Data (FLEED), the Hospital Discharge Register (HDR) of National Institute for Health and Welfare, and the Causes of Death Statistics of Statistics Finland. Individual-level register linkages were conducted using personal identity numbers, which have been assigned to all Finnish residents starting from the year 1969.

FLEED is an annual panel data recording the entire Finnish working-age population. It is constructed from administrative registers of individuals including information on individuals’ labour market status, salaries and other sources of income extracted from tax and other administrative registers, such as from government-run pension registers. FLEED also includes information on relevant individual characteristics and contains general demographic information. FLEED is available from 1988 onwards and for the current study we use the data over the period 1988–2015. HDR contains information on all hospital admissions in Finland, both public and private. Causes of Death Statistics include information on all dates and causes of deaths in Finland. Ethical approval for the study was received from the Research Ethics Committee of National Institute for Health and

Welfare (decision #10/2016§751). Data were linked with the permissions of National Institute for Health and Welfare and Statistics Finland by the appropriate authorities.

Assessment of serious mental disorders

Mental disorder diagnoses have been recorded in HDR according to the ICD-9 with DSM-III-R criteria from 1986 to 1995 and according to the ICD-10 since 1996. We defined mental disorders using the following ICD-10 codes (and their corresponding ICD-9 codes): alcohol abuse (ICD-10: F10), other substance abuse (ICD-10: F11-19), schizophrenia (ICD-10: F20), other non-affective psychoses (ICD-10: F22-29), bipolar disorders (ICD-10: F30-31), depressive disorders (ICD-10: F32-33 and F34.1), other mood disorders (ICD-10: F38-39, F34.0, F34.8 and F34.9) and anxiety disorders (F40-48). All these mental disorders were labelled as serious mental disorders in the current study context where only inpatient data were available. Both primary and secondary diagnoses were used, and an individual could have more than one diagnosis. For sensitivity analyses, a hierarchy of the diagnoses using the following structure was constructed: schizophrenia, other non-affective psychoses, other substance abuse, alcohol abuse, bipolar disorders, depressive disorders, other mood disorders and anxiety disorders. In addition, for those individuals that had information of their biological parents ($n = 1\ 740\ 726$), history of parental mental disorders, that is information whether either or both of the parents were treated for a mental disorder in a hospital, was obtained from the HDR. Parental mental disorders were defined as no mental disorder *vs* any mental disorder in either or both parents (ICD-10 codes: F10 to F48).

Labour market outcomes

All labour market outcomes were obtained from the FLEED. Annual employment status was measured as the employment status during the last week of each year. Individuals who were working either part-time or full-time in a legal employment contract, or self-employed were defined as employed. All others were classified as not employed. Annual earnings were measured as the average of annual wage and salary earnings and self-employed income for each year. Total income was measured as the total taxable income, which includes in addition to annual wage and salary earnings and self-employment income, income transfers (e.g. sick leave allowance), social security benefits (e.g. parental leave benefits) and capital income (capital gains and

dividends). To allow for the comparability of earnings over the years, all income measures were deflated to the base year 2015 using the official consumer price index provided by Statistics Finland. Highest completed secondary or higher education was obtained from the Finnish registry for educational achievement, where it was recorded annually. Individuals who did not have any secondary or higher education and thus had no data in the education registry were categorized as having no secondary or higher education.

Statistical analyses

The associations of serious mental disorders with not being employed and not having any secondary or higher education were examined using analyses where mental disorder diagnoses between ages 15 and 25 (yes/no) was modelled as the exposure and employment or education status between ages 25 and 52 as the outcomes. For these two outcomes, separate models were estimated for all eight different mental disorders and all ages between 25 and 52. All estimates were adjusted for sex, the year of birth and migrant status where a migrant was defined as anyone born outside of Finland (first generation) or born in Finland but with both parents born outside Finland. Additional analyses were conducted, where the associations were adjusted also for parental history of mental disorders. Instead of reporting odds ratios, a relative risk ratio (RR) was considered to be a more interpretable for the present study as the outcomes had a prevalence of more than 10% (36). Hence, a modified Poisson regression was used in the analyses (37, 38). Possible sex differences in the associations were examined by including an interaction term between sex and a mental disorder diagnosis in separate analyses.

Median and mean earnings and total income levels for all ages between ages 25 and 52 were calculated for individuals who were diagnosed with a mental disorder between the age of 15 and 25. In a similar way, mean earnings and total income levels for all ages between ages 25 and 52 were also calculated for those individuals who were not diagnosed with any serious mental disorders. In addition, predicted mean earnings and total income were estimated using linear regression models. Mental disorder diagnoses between ages 15 and 25 (yes/no) were modelled as the exposure and earnings and total income between ages 25 and 52 as the outcomes. The separate models in which sex, the year of birth and migrant status were accounted for were estimated for all eight different mental disorders.

Lastly, the association between mental disorders with employment status and education was also examined using a hierarchy where diagnoses were ranked according to alleged severity, that is from the most (schizophrenia) to least (anxiety disorders) severe. STATA 15.1 (Stata Corp, College Station, TX) was used in all analyses.

Results

In total, 2 055 720 individuals (48.8% women) were included in the analyses. Descriptive statistics of the sample are shown in Table 1, and the number of individuals in these diagnostic groups between ages of 25 and 52 is shown in Table S1. Due to very small number of individuals with other substance abuse diagnosis between ages 15 and 25, and complete follow-up data at the age of 52, results for these analyses are not shown.

The percentage of individuals who were not employed and who did not complete any secondary or higher education between ages 25 and 52 are shown Fig. 1. Overall, the rate of not being in employment was highest among individuals with schizophrenia (range from 89% to 94%). Slightly lower rates were found for individuals diagnosed with other non-affective psychosis (range from 76% to 84%), with alcohol abuse (range from 60% to 70%), with other substance abuse (range

from 63% to 82%) and with bipolar disorders (range from 60% to 75%). Among individuals diagnosed for depressive disorders 52–73% were not employed, and respective rates for other mood disorders were 49–62% and anxiety disorders were 41–59%. Whereas in the comparison population without any of these disorders the rate of not being employed rapidly declined over the follow-up period (i.e. 35% of the individuals were not employed at the age of 25, and 19% of the individuals were not employed at the age of 45), for most mental disorders the rate of not being employed increased over the follow-up period. Exception to this pattern was substance abuse other than alcohol and other mood disorders, where the rate of not being employed also declined over time.

In the comparison population without any mental disorders, 14% to 17% of individuals had not completed some secondary or higher education (Fig. 1). From the different mental disorder groups, individuals with other substance abuse disorder had the highest rate of not completing any secondary or higher education (range from 45% to 70%). Among individuals with alcohol abuse the rate was between 40% and 52%; with schizophrenia between 43% and 54%; and with other non-affective psychosis between 30% and 42%. Relatively similar rates were found for mood disorders and anxiety disorders. For all mental disorders, the rate of not having any secondary or higher education declined over the follow-up period.

The relative risks for mental disorder diagnosis between ages 15 and 25 and not being employed are shown Fig. 2. Overall, schizophrenia was associated with the highest relative risk of not being employed between the ages of 25 and 52. At the age of 25, individuals with schizophrenia had 2.7-fold relative risk of not being employed; at the age of 45, it was over two times higher. A similar trend was also found for other non-affective psychoses. In comparison, alcohol abuse was associated with 2-fold relative risk and other substance abuse with 2.5-fold relative risks of not being employed at the age of 25. At the age of 45, however, these same relative risks were around 1.5 times higher, indicating that the relative risks increased over follow-up. Sex-specific associations indicated that men with alcohol abuse, other substance abuse, schizophrenia and other non-affective psychoses had higher relative risk of not being employed between ages 25 and 40 than women (all *P*-values < 0.01) (Figure S1; Tables S2–S3). After that notable differences were not found (all *P*-values > 0.01).

All mood disorders and anxiety disorders were also associated with higher relative risks of not being employed between the ages of 25 and 52

Table 1. Descriptive statistics of the study population (*n* = 2 055 720)

	<i>N</i> (% of the total number of individuals*)	% women	% migrant	% with parental MD†	Mean age when diagnosed (SD)
Alcohol use disorder	9057 (0.44%)	29.8%	2.9%	32.3%	20.9 (3.0)
Other substance abuse disorders	7432 (0.36%)	31.2%	5.1%	32.4%	21.4 (2.5)
Schizophrenia	6314 (0.31%)	35.9%	4.1%	28.5%	21.4 (2.6)
Other non-affective psychoses	12 604 (0.61%)	42.4%	4.5%	26.6%	21.0 (2.8)
Bipolar disorders	2873 (0.14%)	60.8%	4.0%	29.3%	21.7 (2.6)
Depressive disorders	17 055 (0.83%)	58.0%	4.4%	26.9%	20.5 (2.9)
Other mood disorders	1666 (0.08%)	36.6%	1.9%	24.3%	19.8 (2.2)
Anxiety disorders	21 232 (1.03%)	27.3%	3.6%	22.5%	20.3 (2.3)
No serious mental disorder	1 994 671 (97.03%)	49.1%	12.7%	12.0%	

*Individual could have more than one diagnosis, and thus, the total percentage is higher than 100.

†Total number of individuals with information of both biological parents is 1 740 726.




Fig. 1. Percentage of individuals who were not employed or did not have secondary or higher education between ages 25 and 52 by mental disorder categories.

(Fig. 2). These relative risks were slightly higher for bipolar disorders than for depressive disorders, other mood disorders or anxiety disorders. They also increased up to the age of 40 and then remained rather stable. Sex-specific associations suggested that after the age of 40 the relative risks of not being employed (P -values < 0.01) were slightly higher for women than for men diagnosed any mood or anxiety disorders (Figure S2; Tables S4–S5).

In additional analyses, the history of parental mental disorders was also included as covariate. These results are reported in Table S6. The risk ratios for different mental disorder diagnoses of not being employed and not having any secondary or higher education were only slightly diluted when the history of parental mental disorders was also adjusted.

Figure 3 shows the median earnings and total income levels for the individuals with different mental disorders diagnosed between the age of 15 and 25, and for comparison population. Over half of the individuals with alcohol abuse, other substance abuse, schizophrenia, other non-affective psychoses or bipolar disorders did not have any earnings or

very little earnings between ages 25 and 52. For these groups, the median total income was under or little over 10 000 € between the ages 25 and 52. Whereas the median earnings for individuals with depressive disorders or other mood disorders were also very small, the median total income slightly increased over time. For the anxiety disorders group, the median earnings and total income increased before the age of 35 and decreased after that. These results are in line with the unadjusted mean earnings and total income shown Figure S3. However, the estimated mean earnings and total income, which are shown in Figures S4–S7, suggest that the variation in earnings and total income increases considerably over time.

Figure 4 shows the relative risks for not having any secondary or higher education after the age of 25 for different mental disorder categories. At the age of 25, schizophrenia, other non-affective psychoses, alcohol abuse and other substance abuse were associated between 2.60- to 4.35-fold relative risk of not having completed any secondary or higher education. For other non-affective psychoses, alcohol abuse, and especially for other substance abuse, the relative risks of not having




Fig. 2. Associations of mental disorder diagnosis between ages 15 and 25 with not being employed over the work-life course.

completed any secondary or higher education decreased slightly over follow-up, whereas there was a modest increasing trend among individuals diagnosed with schizophrenia. All mood disorders and anxiety disorders were associated with around 2.5-fold relative risk of not having any secondary or higher education at the age of 25. This relative risk decreased slightly over follow-up among individuals with bipolar disorders. Sex-specific associations showed that the relative risk of not having any secondary or higher education were slightly higher for most of the mental disorders among women (all P -values < 0.01) (Figures S8–S9; Tables S7–S10). The additional analyses, where the history of parental mental disorders was also included as covariate, suggested that the associations were minimally attenuated when the history of parental mental disorders was taken into account (Tables S11).

Sensitivity analyses

The risk ratios for mental disorder diagnosis between ages 15 and 25 with not being employed

and not having any secondary or higher education after the age of 25 using a hierarchy of mental disorder diagnoses are shown in Figures S10 and S11, respectively. Compared with the previously reported associations (Fig. 2), the risk ratios for not being employed were slightly higher for schizophrenia and somewhat diluted for all other mental disorders. Similarly, the risk ratios for not having any secondary or higher education were slightly higher for schizophrenia and slightly lower or very similar for all other mental disorders.

Discussion

The present study examined the associations between serious mental disorders and subsequent employment, earnings, total income and educational outcomes from a work career perspective. We used comprehensive national register data from Finland. Our main finding was that all serious mental disorders were associated with higher rates of not being employed and not having any secondary or higher education, and thus leading to a substantial loss of total earnings and income.




Fig. 3. Median earnings and total income between the age of 25 and 52 for individuals with a mental disorder diagnosis between the ages of 15 and 25.

In our analyses, especially schizophrenia, but also the other non-affective psychoses, alcohol abuse, other substance abuse and bipolar disorders were also associated with considerably higher risk of not being employed and a significant loss of total earnings. Lastly, individuals with mood disorders had considerably low earnings as well. Our income related estimates, however, are conservative since individuals with mental disorders who were not hospitalized between ages 15 to 25 and whose disorder started after age 25 were included in the comparison group, thus lowering the estimates for the labour market losses associated with mental disorders. Moreover, the total income was considerable higher for all mental disorder groups—and especially in schizophrenia—compared with earned income, indicating that income transfers increased considerably the total income of individuals with mental disorders. Lastly, variation in the predicted income increased over time, indicating that some individuals do better than others.

It is well established that indirect costs, such as non-employment, constitute a large part of the total economic burden caused by mental disorders

(3–7). Most previous studies have also shown that most disorders are associated with lower rates of employment and low income (9–20), but not necessarily with low education (28). To our knowledge, the current study is one of the largest to examine the associations of serious mental disorders with subsequent economic outcomes and the first one to compare the associations between different mental disorders that are critical for policy decisions to allocate the healthcare resources to the treatment of these conditions. In addition to showing the associations between mental disorders with lower rates of employment, earnings, total income and education, the results showed that the effect sizes of these associations—except for individuals diagnosed with drug-related substance abuse disorder—increased over time, indicating that for individuals diagnosed with a serious mental disorder before the age of 25 the economic prospects are worse over the long run. Although this might not lower their quality of life, pensions in Finland and in other Nordic countries are tightly linked to earnings before the official retirement age, thus persons with mental disorders will suffer from a low




Fig. 4. Associations of mental disorder diagnosis between ages 15 and 25 with no secondary or higher education between ages 25 and 52.

income also after retiring. Moreover, although Finland has a universal health care and welfare system in which coverage is intended to provide economic security during unemployment and to compensate the income losses during periods of illness, it is not known how well these policies help individuals to obtain work after an onset of mental disorders. Although our study did not address specific policies, our findings could be used to project how well policies in different countries for individuals with mental disorders work.

Plausible mechanisms

There are several plausible explanations for the differences we found between mental disorders. Individuals who have early and frequent episodes of acute mental health problems are likely to need more time off from work or studies, which in turn is likely to result in their exit from the labour market and widen the income differential. Individuals with serious mental disorders, such as schizophrenia and bipolar disorders, have difficulties in mental operations that facilitate social interactions

(39), including emotion recognition and expression. These difficulties are likely to lead to problems in workplaces where expressing and responding to emotions is required. In addition, individuals with mental disorders are subject to stigmatization, studies showing that individuals with serious mental disorders experience discrimination in multiple domains of life (40). Whereas stigmatization has been shown to lead to delayed access to care (41, 42), it could also lead to the loss of a job and difficulties in finding or even applying for a new one. Mental disorders have also been shown to reduce work performance (43), which may substantially reduce earnings in the long run, especially at workplaces where earnings are based on performance evaluations and ratings at work. Whereas cognitive deficits are a central feature of psychotic disorders and especially schizophrenia (44), they are less present—with the exception of chronic forms—in other mental disorders such as depression (45). Although good general cognitive capacity has been associated with higher earnings and better career prospects (46, 47), these effects are rather modest, and there are also many jobs in

which high cognitive capacity is not required, and thus individuals with difficulties in cognitive performance can be employed to these jobs. Overall, schizophrenia has been considered more chronic and more disabling than many other disorders. The present results support this conclusion. Last, it must be noted that the association between mental disorders with employment and education is reciprocal and, likely, quite complex as unemployment and low education have been consistently associated with increased risk of mental disorders (31–33).

Strengths and limitations

The use of register data has obvious strengths. All legal employment contracts in Finland are recorded in the data. Although we measured employment status only during the last week of each year, the annual earnings measure that we use covers all earnings and total income during each year, even from the shortest part-time and temporary contracts. This register-based information, linked to HDR, enabled us to examine the long-term association of serious mental disorders with employment and educational outcomes using nationally representative data.

This study has some important limitations. First, we used register data that were not originally designed for research purposes. Although the diagnostic validity of schizophrenia (48) and bipolar disorder I (49) diagnoses in the Finnish register data have been reported to be good, we did not have diagnostic validity information for other mental disorders. Second, we did not have data on the severity of mental disorders. As our data contained only individuals who were diagnosed with mental disorders in Finnish hospitals, individuals with more severe symptoms were included in the data. Individuals who did not seek treatment or who received treatment in primary care, in outpatient clinics or in voluntary/community sector were not included. Moreover, individuals from a high-income background could be more likely to use private care services, which are also available in Finland to a limited extent. Due to these reasons, current findings describe the labour market outcomes for those individuals who have been ill-enough to have been hospitalized for mental disorders during their young adulthood. Thus, present results may not be applicable to individuals with less severe or well-managed serious mental disorders. Third, selective survival bias, that is selective attrition where individuals with more severe symptoms (50, 51) and from worse socioeconomic backgrounds are more likely to die, could explain part

of the findings. Fourth, due to differences in welfare and healthcare policies between countries, generalization of the current study results outside the Finnish setting might be limited.

Conclusion

To conclude, the results of the present study using administrative register data from Finland suggest that individuals who have been hospitalized with substance abuse, psychosis and mood or anxiety related mental disorders have an increased risk of not being employed and not having any secondary or higher education over the work-life course. As a consequence of this, serious mental disorders lead to substantial losses of earnings and total income.

Acknowledgements

This work was supported by the Academy of Finland (310591 to C.H.) and Palkansaajasäätiö Foundation.

Conflict of interests

None.

Author contribution

CH, ME and PB contributed to the concept and design of the study. CH conducted the data analysis and wrote the first draft of the manuscript. MA, SL and PB helped to conduct the data analysis. All authors contributed to the interpretation of the results, manuscript revision and approved the final version of the manuscript. CH is the guarantor. The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

Data availability statement

The data that support the findings of this study are available from the National Institute for Health and Welfare and Statistics Finland. Restrictions apply to the availability of these data, which were used under license for this study. For information on accessing the data see www.stat.fi and www.thl.fi.

References

1. WHITEFORD H, DEGENHARDT L, REHM J et al. Global burden of disease attributable to mental and substance use disorders: findings from the Global Burden of Disease Study 2010. *Lancet* 2013;**382**:1575–1586.
2. ABAJOBIR AA, ABATE KH, ABBAFATI C et al. Global, regional, and national disability-adjusted life-years (DALYs)

- for 333 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017;**390**:1260–1344.
3. BLOOM DE, CAFIERO E, JANÉ-LLOPIS E et al. The global economic burden of noncommunicable diseases. PGDA Working Papers 8712, Program on the Global Demography of Aging.
 4. GREENBERG PE, FOURNIER A-A, SISITSKY T, PIKE CT, KESSLER RC. The economic burden of adults with major depressive disorder in the United States (2005 and 2010). *J Clin Psychiatry* 2015;**76**:155–162.
 5. CLOUTIER M, GREENE M, GUERIN A, TOUYA M, WU E. The economic burden of bipolar I disorder in the United States in 2015. *J Affect Disord* 2018;**226**:45–51.
 6. CHONG HY, TEOH SL, WU DB-C et al. Global economic burden of schizophrenia: a systematic review. *Neuropsychiatr Dis Treat* 2016;**12**:357–373.
 7. CLOUTIER M, SANON AIGBOGUN M, GUERIN A et al. The economic burden of Schizophrenia in the United States in 2013. *J Clin Psychiatry* 2016;**77**:764–771.
 8. Schizophrenia Commission. The Abandoned Illness: A Report from the Schizophrenia Commission. London 2012. <http://www.schizophreniacommission.org.uk/the-report/>
 9. ZIMMERMAN FJ, KATON W. Socioeconomic status, depression disparities, and financial strain: What lies behind the income-depression relationship? *Health Econ* 2005;**14**:1197–1215.
 10. WHOOLEY M, KIEFE C, CHESNEY M, MARKOVITZ J, MATTHEWS K, HULLEY S. Depressive symptoms, unemployment, and loss of income. *Arch Intern Med* 2002;**162**:2614–2620.
 11. SAMPSON N, KESSLER RC, SUSUKIDA R et al. Long-term effects of mental disorders on employment in the National Comorbidity Survey ten-year follow-up. *Soc Psychiatry Psychiatr Epidemiol* 2015;**50**:1657–1668.
 12. LERNER D, HENKE RM. What does research tell us about depression, job performance, and work productivity? *J Occup Environ Med* 2008;**50**:401–410.
 13. HAKULINEN C, ELOVAINIO M, PULKKI-RÄBACK L et al. Depressive symptoms and long-term income: The Young Finns Study. *J Affect Disord* 2016;**204**:120–123.
 14. AGERBO E, BYRNE M, EATON WW, MORTENSEN PB. Marital and labor market status in the long run in schizophrenia. *Arch Gen Psychiatry* 2004;**61**:28–33.
 15. FALK J, BURSTRÖM B, DALMAN C, JÖRGENSEN L, BRUCE D, NYLÉN L. Employment and income among first-time cases diagnosed with non-affective psychosis in Stockholm, Sweden: a follow-up study 2004/2005–2010. *Soc Psychiatry Psychiatr Epidemiol* 2016;**51**:259–267.
 16. DAVIDSON M, KAPARA O, GOLDBERG S, YOFFE R, NOY S, WEISER M. A nation-wide study on the percentage of schizophrenia and bipolar disorder patients who earn minimum wage or above. *Schizophr Bull* 2016;**42**:443–447.
 17. BANERJEE S, CHATTERJI P, LAHIRI K. Effects of psychiatric disorders on labor market outcomes: a latent variable approach using multiple clinical indicators. *Health Econ* 2015;**19**:n/a–n/a.
 18. ETTNER SL, FRANK RG, KESSLER RC. The impact of psychiatric disorders on labor market outcomes. *Ind Labor Relations Rev* 1997;**51**:64–81.
 19. JØRGENSEN MB, PEDERSEN J, THYGESEN LC et al. Alcohol consumption and labour market participation: a prospective cohort study of transitions between work, unemployment, sickness absence, and social benefits. *Eur J Epidemiol* 2019;**34**:397–407.
 20. SAREEN J, AFIFI TO, McMILLAN KA, ASMUNDSON GJG. Relationship between household income and mental disorders. *Arch Gen Psychiatry* 2011;**68**:419.
 21. KESSLER RC, HEERINGA S, LAKOMA M et al. Individual and societal effects of mental disorders on earnings in the United States: results from the National Comorbidity Survey Replication. *Am J Psychiatry* 2008;**165**:703–711.
 22. DOHRENWEND BP, LEVAV I, SHROUT PE et al. Socioeconomic status and psychiatric disorders: the causation-selection issue. *Science* 1992;**255**:946–952.
 23. MARWAHA S, JOHNSON S, BEBBINGTON P, STAFFORD M, ANGERMEYER MC. Rates and correlates of employment in people with schizophrenia in the UK, France and Germany. *Br J Psychiatry* 2007;**191**:30–37.
 24. MARWAHA S, JOHNSON S. Schizophrenia and employment: A review. *Soc Psychiatry Psychiatr Epidemiol* 2004;**39**:337–349.
 25. MARWAHA S, DURRANI A, SINGH S. Employment outcomes in people with bipolar disorder: a systematic review. *Acta Psychiatr Scand* 2013;**128**:179–193.
 26. LEE S, TSANG A, BRESLAU J et al. Mental disorders and termination of education in high-income and low- and middle-income countries: epidemiological study. *Br J Psychiatry* 2009;**194**:411–417.
 27. MOJTABAI R, SAMPSON N, HWANG I, EATON WW, KESSLER RC, STUART EA. Long-term effects of mental disorders on educational attainment in the National Comorbidity Survey ten-year follow-up. *Soc Psychiatry Psychiatr Epidemiol* 2015;**50**:1577–1591.
 28. TEMPELAAR WM, TERMORSHUIZEN F, MACCABE JH, BOKS MPM, KAHN RS. Educational achievement in psychiatric patients and their siblings: A register-based study in 30 000 individuals in the Netherlands. *Psychol Med* 2017;**47**:776–784.
 29. STRINGHINI S, CARMELI C, JOKELA M et al. Socioeconomic status and the 25 × 25 risk factors as determinants of premature mortality: a multicohort study and meta-analysis of 1.7 million men and women. *Lancet* 2017;**389**:1229–1237.
 30. STRINGHINI S, CARMELI C, JOKELA M et al. Socioeconomic status, non-communicable disease risk factors, and walking speed in older adults: multi-cohort population based study. *BMJ* 2018;**360**:k1046.
 31. PAUL KI, MOSER K. Unemployment impairs mental health: Meta-analyses. *J Vocat Behav* 2009;**74**:264–282.
 32. LORANT V, DELIEGE D, EATON W, ROBERT A, PHILIPPOT P, ANSSEAU M. Socioeconomic inequalities in depression: a meta-analysis. *Am J Epidemiol* 2003;**157**:98–112.
 33. MUNTANER C, EATON WW, MIECH R, O'CAMPO P. Socioeconomic position and major mental disorders. *Epidemiol Rev* 2004;**26**:53–62.
 34. KESSLER RC, BERGLUND P, DEMLER O, JIN R, WALTERS EE. Lifetime prevalence and age-of-onset distributions' of DSM-IV disorders in the national comorbidity survey replication. *Arch Gen Psychiatry* 2005;**62**:593–602.
 35. PEDERSEN CB, MORS O, BERTELSEN A et al. A comprehensive nationwide study of the incidence rate and lifetime risk for treated mental disorders. *JAMA Psychiatry* 2014;**71**:573–581.
 36. SCHMIDT CO, KOHLMANN T. When to use the odds ratio or the relative risk? *Int J Public Health* 2008;**53**:165–167.
 37. ZOU G. A modified Poisson regression approach to prospective studies with binary data. *Am J Epidemiol* 2004;**159**:702–706.
 38. CUMMINGS P. Methods for estimating adjusted risk ratios. *Stata J* 2009;**9**:175–196.
 39. COTTER J, GRANGER K, BACKX R, HOBBS M, LOOI CY, BARNETT JH. Social cognitive dysfunction as a clinical marker: A

- systematic review of meta-analyses across 30 clinical conditions. *Neurosci Biobehav Rev* 2018;**84**:92–99.
40. THORNICROFT G, BROHAN E, ROSE D, SARTORIUS N, LEESE M. Global pattern of experienced and anticipated discrimination against people with schizophrenia: a cross-sectional survey. *Lancet* 2009;**373**:408–415.
 41. GRONHOLM PC, THORNICROFT G, LAURENS KR, EVANS-LACKO S. Mental health-related stigma and pathways to care for people at risk of psychotic disorders or experiencing first-episode psychosis: A systematic review. *Psychol Med* 2017;**47**:1867–1879.
 42. KULAR A, PERRY BI, BROWN L et al. Stigma and access to care in first-episode psychosis. *Early Interv Psychiatry* 2018;1–6.
 43. STEWART WF, RICCI JA, CHEE et al. Cost of lost productive work time among US workers with depression. *JAMA* 2003;**289**:3135–3144.
 44. SCHAEFER J, GIANGRANDE E, WEINBERGER DR, DICKINSON D. The global cognitive impairment in schizophrenia: Consistent over decades and around the world. *Schizophr Res* 2013;**150**:42–50.
 45. ROCK PL, ROISER JP, RIEDEL WJ, BLACKWELL AD. Cognitive impairment in depression: a systematic review and meta-analysis. *Psychol Med* 2014;**44**:2029–2040.
 46. STRENZE T. Intelligence and socioeconomic success: A meta-analytic review of longitudinal research. *Intelligence* 2007;**35**:401–426.
 47. MURNANE RJ, WILLETT JB, DUHALDEBORDE Y, TYLER JH. How important are the cognitive skills of teenagers in predicting subsequent earnings? *Policy Anal* 2012;**19**:547–568.
 48. MÄKIKYRÖ T, ISOHANNI M, MORING J, HAKKO H, HOVATTA I, LÖNNQVIST J. Accuracy of register-based schizophrenia diagnoses in a genetic study. *Eur Psychiatry* 1998;**13**:57–62.
 49. KIESEPPA T, PARTONEN T, KAPRIO J, LÖNNQVIST J. Accuracy of register- and record-based bipolar I disorder diagnoses in Finland; a study of twins. *Acta Neuropsychiatr* 2000;**12**:106–109.
 50. LUMME S, PIKKOLA S, MANDERBACKA K, KESKIMÄKI I. Excess mortality in patients with severe mental disorders in 1996–2010 in Finland. *PLoS ONE* 2016;**11**:e0152223.
 51. SIMON GE, STEWART C, YARBOROUGH BJ et al. Mortality rates after the first diagnosis of psychotic disorder in adolescents and young adults. *JAMA Psychiatry* 2018;**75**:254–260.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Figure S1. Sex-specific associations between alcohol abuse, other substance abuse, schizophrenia, and other non-affective psychosis with not being employed over the work life course.

Figure S2. Sex-specific associations between bipolar disorders, depressive disorders, other mood disorders, and anxiety disorders with not being employed over the work life course.

Figure S3. Mean earnings and total income between the age of 25 and 52 for individuals with a mental disorder diagnosis between the ages of 15 and 25.

Figure S4. Predicted total earnings for individuals with alcohol abuse, other substance abuse, schizophrenia, and other non-affective psychosis over the work life course.

Figure S5. Predicted total earnings for individuals with bipolar disorders, depressive disorders, other mood disorders, and anxiety disorders over the work life course.

Figure S6. Predicted total income for individuals with alcohol abuse, other substance abuse, schizophrenia, and other non-affective psychosis over the work life course.

Figure S7. Predicted total income for individuals with bipolar disorders, depressive disorders, other mood disorders, and anxiety disorders over the work life course.

Figure S8. Sex-specific associations between alcohol abuse, other substance abuse, schizophrenia, and other non-affective psychosis with not having completed any secondary or higher education over the work life course.

Figure S9. Sex-specific associations between bipolar disorders, depressive disorders, other mood disorders, and anxiety disorders with not having completed any secondary or higher education over the work life course.

Figure S10. Associations of mental disorder diagnosis (hierarchical model) between ages 15 and 25 with not being employed between ages 25 and 52.

Figure S11. Associations of mental disorder diagnosis (hierarchical model) between ages 15 and 25 with no secondary or higher education between ages 25 and 52.

Table S1. Number of individuals in the study population.

Table S2. Sex-specific associations between alcohol abuse and other substance abuse with not being employed over the work life course.

Table S3. Sex-specific associations between schizophrenia and other non-affective psychosis with not being employed over the work life course.

Table S4. Sex-specific associations between bipolar disorders and depressive disorders with not being employed over the work life course.

Table S5. Sex-specific associations between other mood disorders and anxiety disorders with not being employed over the work life course.

Table S6. The associations between mental disorders with not being employed at the age of 25, 30, 40, and 50.

Table S7. Sex-specific associations between alcohol abuse and other substance abuse with not having any secondary or higher education over the work life course.

Table S8. Sex-specific associations between schizophrenia and other non-affective psychosis with not having any secondary or higher education over the work life course.

Table S9. Sex-specific associations between bipolar disorders and depressive disorders with not having any secondary or higher education over the work life course.

Table S10. Sex-specific associations between other mood disorders and anxiety disorders with not having any secondary or higher education over the work life course.

Table S11. The associations between mental disorders with no higher or secondary education at the age of 25, 30, 40, and 50.